

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.



DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF HOUSTON

HOUSTON, TEXAS

NASA CR-

160266

(NASA-CR-160266) A MODIFICATION OF THE  
LIKELIHOOD EQUATIONS FOR NORMAL CLASSES WITH  
FIELD STRUCTURE, PRELIMINARY REPORT (Houston  
Univ.) 10 p HC A02/MP A01 CSCL 12A

N79-27894

Unclas  
29202

G3/64

A MODIFICATION OF THE LIKELIHOOD  
EQUATIONS FOR NORMAL CLASSES  
WITH FIELD STRUCTURE  
PRELIMINARY REPORT

BY  
CHARLES PETERS  
REPORT 73 JUNE 1979

PREPARED FOR  
EARTH OBSERVATION DIVISION, JSC  
UNDER  
CONTRACT NAS-9-15543



HOUSTON, TEXAS 77004

A MODIFICATION OF THE LIKELIHOOD  
EQUATIONS FOR NORMAL CLASSES WITH  
FIELD STRUCTURE. (PRELIMINARY REPORT)

by

Charles Peters  
Department of Mathematics  
University of Houston  
Houston, Texas

Report #73  
NAS-9-15543 Mod 2S  
June 1979

## 1. Introduction

A possible objection to the use of UHMLE or CLASSY in conjunction with AMOEBA is that both these algorithms ignore the association of pixels in fields. Indeed AMOEBA is based on the explicit assumption that pixels in the same field represent the same real class [1], while the assumptions underlying the maximum likelihood algorithms imply that the classification of a pixel is independent of the classification of other pixels. In this report a statistical model based on normal mixtures is proposed which takes into account the organization of LANDSAT agricultural data into fields which are homogeneous as to crop type. Likelihood equations for the parameters of the model are derived which may be solved iteratively as in UHMLE.

## 2. The Model

We assume that the data elements (pixel data vectors) are real  $n$ -vectors each from one of the statistical populations  $\Pi_1, \dots, \Pi_m$  with  $n$ -variate density functions  $p(x|\Pi_\ell)$ ,  $\ell = 1, \dots, m$ . We assume that the data is organized into sets (fields)  $F_1, \dots, F_p$ , where  $F_j$  has  $N_j$  data elements which have been previously ordered in some arbitrary fashion so that the data elements in  $F_j$  form a  $nN_j$ -dimensional vector denoted by  $x_j = \begin{matrix} x_{j1} \\ \vdots \\ x_{jN_j} \end{matrix}$

Define random variables  $\{\theta_{jk} \in \{1, \dots, m\} | j=1, \dots, p; k=1; \dots, N_j\}$  by  $\theta_{jk}=\ell$  if and only if  $x_{jk}$  is from  $\Pi_\ell$ . We assume that all the observations from  $F_j$  are

from the same class, so that we may write  $\theta_{jk} = \theta_j$  for all  $j = 1, \dots, p; k, \ell = 1, \dots, N_j$ . Finally, we assume that  $(x_1, \theta_1), \dots, (x_p, \theta_p)$  are independent, that the  $\theta_j$ 's are identically distributed, that

$\alpha_\ell = \text{Prob} [\theta_j = \ell] > 0$  and that  $\sum_{\ell=1}^m \alpha_\ell = 1$ . Under the stated assumptions,

the joint density of  $x_1, \dots, x_p$  is  $p(x_1, \dots, x_p) = \prod_{j=1}^p \sum_{\ell=1}^m \alpha_\ell p(x_j)$ ,

where  $p_\ell(x_j) = p(x_{j1}, \dots, x_{jN_j} | \theta_j = \ell)$  is the joint density of the elements of  $F_j$  given that  $F_j$  represents class  $\Pi_\ell$ .

Let  $N = N_1 + \dots + N_p$  and for each  $\ell$  let  $M_\ell$  denote the total number of the  $N$  observations  $x_{jk}$  which come from class  $\Pi_\ell$ . The following proposition shows that with reasonable restrictions on the field sized  $N_j$  the values of  $\{M_\ell: \ell = 1, \dots, m\}$  can be inferred from a knowledge of the parameters  $\alpha_\ell$ . Thus, acreage estimates of the classes can be derived from estimates of the parameters  $\alpha_\ell$ .

Proposition 1: (a)  $E(M_\ell) = \alpha_\ell N$

(b)  $\frac{M_\ell}{N} \rightarrow \alpha_\ell$  in probability as  $p \rightarrow \infty$  if and only if

$$\lim_{p \rightarrow \infty} \frac{1}{N^2} \sum_{j=1}^p N_j^2 = 0.$$

(c) If  $\sum_{j=1}^{\infty} \frac{N_j^2}{j^2} < \infty$ , then  $\frac{M_\ell}{N} \rightarrow \alpha_\ell$  almost surely.

Proof: (a) Write  $M_\ell = \sum_{j=1}^p \sum_{k=1}^{N_j} x_{\ell}(\theta_{jk})$

$$= \sum_{j=1}^p N_j x_{\ell}(\theta_j)$$

$$\text{where } x_\ell(r) = \begin{cases} 1 & r = \ell \\ 0 & r \neq \ell \end{cases}$$

$$\text{Then } E(M_\ell) = \sum_{j=1}^p N_j E(x_\ell(\theta_j)) = \sum_{j=1}^p N_j \alpha_\ell = N\alpha_\ell.$$

(b) Since  $\frac{M_\ell}{N} - \alpha_\ell = \frac{M_\ell}{N} - \left(E \frac{M_\ell}{N}\right)$  is bounded, it converges to zero in probability iff  $\text{var } \frac{M_\ell}{N} \rightarrow 0$  as  $p \rightarrow \infty$ . Since the terms  $N_j x_\ell(\theta_j)$  are independent,

$$\text{var} \left( \frac{M_\ell}{N} \right) = \frac{1}{N^2} \sum_{j=1}^p N_j^2 \text{var} (x_\ell(\theta_j)) = \frac{1}{N^2} \sum_{j=1}^p N_j^2 \alpha_\ell (1 - \alpha_\ell).$$

The conclusion follows.

(c) The assertion follows immediately from Kolmogorov's version of the strong law of large numbers [3].

### 3. Maximum Likelihood Estimation of the Parameters

In this section we suppose that the class conditional densities  $p(x|\Pi_\ell)$  of the data elements  $x_{jk}$  are n-variate normal  $N(x; \mu_\ell, \Sigma_\ell)$  and that  $\{x_{jk}: k=1, \dots, N_1\}$  are class conditionally independent; i.e., that

$$p_\ell(x_j) = \prod_{k=1}^{N_j} N(x_{jk}; \mu_\ell, \Sigma_\ell).$$

for  $j=1, \dots, p$ . In this case the joint density of  $x_1, \dots, x_p$ ,

$$p(x_1, \dots, x_p) = \prod_{j=1}^p \sum_{\ell=1}^m \alpha_{\ell} \prod_{k=1}^{N_j} N(x_{jk}; \mu_{\ell}, \Sigma_{\ell}),$$

is parametrized by  $\{(\alpha_{\ell}, \mu_{\ell}, \Sigma_{\ell}) | \ell=1, \dots, m\}$  where  $\alpha_{\ell} \geq 0$ ,  $\sum \alpha_{\ell} = 1$ ,  $\mu_{\ell} \in \mathbb{R}^n$ , and  $\Sigma_{\ell}$  is a real  $n \times n$  positive definite symmetric matrix. Whenever a density is evaluated using estimates of its parameters, we denote it, e.g., by  $\hat{p}(x_1, \dots, x_p)$ . By a maximum likelihood estimate (MLE) of the parameters  $\{(\alpha_{\ell}, \mu_{\ell}, \Sigma_{\ell})\}$  we mean an element  $\{(\hat{\alpha}_{\ell}, \hat{\mu}_{\ell}, \hat{\Sigma}_{\ell}) | \ell=1, \dots, m\}$  of the parameter set which locally maximizes  $\hat{p}(x_1, \dots, x_p)$ . By arguments similar to those used in [2], the following necessary conditions for a MLE are derived.

$$1) \quad \frac{1}{p} \sum_{j=1}^p \frac{\hat{p}_{\ell}(x_j)}{\hat{p}(x_j)} \leq 1 \quad \text{with equality when } \alpha_{\ell} > 0$$

$$2) \quad \hat{\mu}_{\ell} = \frac{\sum_{j=1}^p \frac{N_j \hat{p}_{\ell}(x_j)}{\hat{p}(x_j)} \bar{x}_j}{\sum_{j=1}^p \frac{N_j \hat{p}_{\ell}(x_j)}{\hat{p}(x_j)}}$$

$$3) \quad \hat{\Sigma}_{\ell} = \frac{\sum_{j=1}^p \frac{\hat{p}_{\ell}(x_j)}{\hat{p}(x_j)} \sum_{k=1}^{N_j} (x_{jk} - \hat{\mu}_{\ell})(x_{jk} - \hat{\mu}_{\ell})^T}{\sum_{j=1}^p \frac{N_j \hat{p}_{\ell}(x_j)}{\hat{p}(x_j)}}$$

In equation (2)  $\bar{x}_j = \frac{1}{N_j} \sum_{k=1}^{N_j} x_{jk}$  is the mean of the  $j$ th field observations.

By multiplying (1) by  $\hat{\alpha}_{\ell}$  we obtain

$$4) \quad \hat{\alpha}_{\ell} = \frac{1}{p} \sum_{j=1}^p \frac{\alpha_{\ell} \hat{p}_{\ell}(x_j)}{\hat{p}(x_j)}$$



which, together with (2) and (3) suggests an iterative procedure for solution of the likelihood equations (2) - (4) analogous to that used in UHMLE [2]. However, the likelihood equations can be considerably simplified by observing that the sequence  $(\bar{X}_1, S_1), \dots, (\bar{X}_p, S_p)$ , is a sufficient statistic for the model, where  $S_j$  is the sample scatter matrix of the  $j$ th field:

$$S_j = \sum_{k=1}^{N_j} (x_{jk} - \bar{X}_j) (x_{jk} - \bar{X}_j)^T.$$

Equation (3) may be rewritten

$$\begin{aligned} 5) \quad \hat{\Sigma}_\ell = & \frac{p}{\sum_{j=1}^p} \frac{\hat{p}_\ell(x_j)}{\hat{p}(x_j)} S_j \bigg/ \frac{p}{\sum_{j=1}^p} \frac{N_j \hat{p}_\ell(x_j)}{\hat{p}(x_j)} \\ & + \frac{p}{\sum_{j=1}^p} \frac{N_j \hat{p}_\ell(x_j)}{\hat{p}(x_j)} (\bar{X}_j - \hat{\mu}_\ell) (\bar{X}_j - \hat{\mu}_\ell)^T \bigg/ \frac{p}{\sum_{j=1}^p} \frac{N_j \hat{p}_\ell(x_j)}{\hat{p}(x_j)} \end{aligned}$$

The sufficiency of  $\{\bar{X}_j, S_j\}_{j=1}^p$  implies that

$$\frac{\hat{p}_\ell(x_j)}{\hat{p}(x_j)} = \frac{\hat{q}_\ell(\bar{X}_j, S_j)}{\hat{q}(\bar{X}_j, S_j)}$$

where  $\hat{q}_\ell(\bar{X}_j, S_j)$  is the estimated joint density of  $\bar{X}_j$  and  $S_j$  given that  $F_j$  represents class  $\ell$  and  $\hat{q}(\bar{X}_j, S_j) = \sum_{\ell=1}^m \hat{\alpha}_\ell \hat{q}_\ell(\bar{X}_j, S_j)$ . The joint density



$\hat{q}_\ell(\bar{x}_j, S_j)$  may be expressed as

$$q_\ell(\bar{x}_j, S_j) = N_n(\bar{x}_j; \hat{\mu}_\ell, \frac{1}{N_j} \hat{\Sigma}_\ell) W_n(S_j; N_j-1, \hat{\Sigma}_\ell)$$

where  $N_n(\bar{x}_j; \hat{\mu}_\ell, \frac{1}{N_j} \hat{\Sigma}_\ell)$  is the n-variate normal density of  $\bar{x}_j$  and  $W_n(S_j; N_j-1, \hat{\Sigma}_\ell)$  is the Wishart density of  $S_j$  with  $N_j-1$  degrees of freedom [3].

Thus the likelihood equations may be written as

$$6) \quad \hat{\alpha}_\ell = \frac{1}{p} \sum_{j=1}^p \frac{\hat{\alpha}_\ell \hat{q}_\ell(\bar{x}_j, S_j)}{\hat{q}_\ell(\bar{x}_j, S_j)}$$

$$7) \quad \hat{\mu}_\ell = \frac{\sum_{j=1}^p \frac{N_j \hat{q}_\ell(\bar{x}_j, S_j)}{\hat{q}_\ell(\bar{x}_j, S_j)} \bar{x}_j}{\sum_{j=1}^p \frac{N_j \hat{q}_\ell(\bar{x}_j, S_j)}{\hat{q}_\ell(\bar{x}_j, S_j)}}$$

$$8) \quad \hat{\Sigma}_\ell = \frac{\sum_{j=1}^p \frac{\hat{q}_\ell(\bar{x}_j, S_j)}{\hat{q}_\ell(\bar{x}_j, S_j)} S_j}{\sum_{j=1}^p \frac{N_j \hat{q}_\ell(\bar{x}_j, S_j)}{\hat{q}_\ell(\bar{x}_j, S_j)}} + \frac{\sum_{j=1}^p \frac{N_j \hat{q}_\ell(\bar{x}_j, S_j)}{\hat{q}_\ell(\bar{x}_j, S_j)} (\bar{x}_j - \hat{\mu}_\ell)(\bar{x}_j - \hat{\mu}_\ell)^T}{\sum_{j=1}^p \frac{N_j \hat{q}_\ell(\bar{x}_j, S_j)}{\hat{q}_\ell(\bar{x}_j, S_j)}}$$

Equations (6) - (8) are to be used as the basis of the iteration procedure.

Indeed when each  $N_j = 1$  they reduce to the likelihood equations employed in UHMLE.

#### 4. Concluding Remarks.

The questions of the existence of a consistent MLE as  $p \rightarrow \infty$  and the

local convergence of the iterative procedure will be addressed in a future report. We remark that the standard consistency results of Cramer, Chanda, and Wald (see [2] for references) are not directly applicable since the  $(\bar{x}_j, S_j)$  are not identically distributed. Numerical results will also be reported at a later date.

## REFERENCES

1. J. Bryant, "On the clustering of multidimensional pictorial data", to appear in Pattern Recognition.
2. B. C. Peters, Jr. and H. F. Walker, "An iterative procedure for obtaining maximum likelihood estimates of the parameters for a mixture of normal distributions", SIAM J. Appl. Math., B, 35, 2, Sept. 1978.
3. S. S. Wilks, Mathematical Statistics, Wiley and Sons, New York, 1962.